# Physics 2414, Spring 2005 Group Exercise 2, Feb 3, 2005

Name 1:		OUID 1: .	
Name 2:		OUID 2:	
Name 3:		OUID 3:	
Name 4:		OUID 4:	
Section Number:			

# **Frictional Force**

#### Notation

 $\vec{\mathbf{F}}_f$  - Frictional force (static or kinetic).

 $\vec{\mathbf{F}}_s$  - Static frictional force.

 $\vec{\mathbf{F}}_k$  - Kinetic frictional force.

## Description

A tank (of mass 10 kg with water of mass 90 kg) open to the sunlight (so that the water evaporates) is attached to a mass  $m_0 = 50$  kg using a massless frictionless pulley as shown in figure 1. The surface on which the tank rests has coefficient of static friction  $\mu_s = 0.8$ , and coefficient of kinetic friction  $\mu_k = 0.5$ .





Figure 1: Diagram showing the water#tank being pulled by mass  $m_0$ .

Due to evaporation, the weight (=Mg) of the tank+water varies as shown in the plot 1.



Plot 1: Mg verses time


## Problems

- 1. Normal force:
- (a) Draw the free body diagram for the system consisting of the tank+water.



(b) Write the expression for the normal force on the tank+water system.

(c) Plot the normal force as a function of time in plot 2.




Plot 2: N verses time

#### 2. Friction:

(a) Write the force equation for the tank+water in the horizontal direction.

(b) At what time does the tank start to move? (Hint:  $T = m_0 g = 490$  N before the tank starts moving; the acceleration of the tank is zero before it starts to move; and  $F_f = \mu_s N$  instantly before the tank starts to move.)

(c) Plot the frictional force as a function of time in plot 3. (Hint:  $F_s \le \mu_s N$  and  $F_k = \mu_k N$ .)



Plot 3:  $F_f$  verses time

(d) What can you tell about the acceleration of the tank after it starts to move? Does the acceleration of the tank attain a constant value after all the water in the tank evaporates? Give a qualitative answer.