

Exam No. 02 (Fall 2013)

PHYS 320: Electricity and Magnetism I

Date: 2013 Oct 23

1. **(20 points.)** The electric field due to a point dipole \mathbf{d} at a distance \mathbf{r} away from dipole is given by the expression

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{1}{r^3} [3(\mathbf{d} \cdot \hat{\mathbf{r}})\hat{\mathbf{r}} - \mathbf{d}]. \quad (1)$$

Consider the case when the point dipole is positioned at the origin and is pointing in the z -direction, i.e., $\mathbf{d} = d\hat{\mathbf{z}}$.

- (a) Qualitatively plot the electric field lines for the dipole \mathbf{d} . (Hint: You do not have to depend on Eq. (1) for this purpose. An intuitive knowledge of electric field lines should be the guide.)
- (b) Find the (simplified) expression for the electric field on the positive z -axis. (Hint: On the positive z -axis we have, $\hat{\mathbf{r}} = \hat{\mathbf{z}}$ and $r = z$.)
2. **(30 points.)** Consider a solid sphere of radius R with total charge Q distributed inside the sphere with a charge density

$$\rho(\mathbf{r}) = br^2 \theta(R - r), \quad (2)$$

where r is the distance from the center of sphere, and $\theta(x) = 1$, if $x > 0$, and 0 otherwise.

- (a) Integrating the charge density over all space gives you the total charge Q . Thus, determine the constant b in terms of Q and R .
- (b) Using Gauss's law find the electric field inside and outside the sphere.
- (c) Plot the electric field as a function of r .
3. **(20 points.)** In class we evaluated the electric potential due to a solid sphere with uniform charge density Q . The angular integral in this evaluation involved the integral

$$\frac{1}{2} \int_{-1}^1 dt \frac{1}{\sqrt{r^2 + r'^2 - 2rr't}}. \quad (3)$$

Evaluate the integral for $r < r'$ and $r' < r$, where r and r' are distances measured from the center of the sphere. (Hint: Substitute $r^2 + r'^2 - 2rr't = y$.)

4. **(30 points.)** The charge density for a point charge q_a is described by

$$\rho(\mathbf{r}) = q_a \delta^{(3)}(\mathbf{r} - \mathbf{r}_a), \quad (4)$$

where \mathbf{r}_a is the position of the charge.

- (a) Evaluate the electric potential due to the point charge using

$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int d^3r' \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}. \quad (5)$$

(Hint: Use the δ -function property to evaluate the integrals.)

- (b) Evaluate the electric field due to the point charge by finding the gradient of the electric potential you calculated using Eq. (5),

$$\mathbf{E}(\mathbf{r}) = -\nabla\phi(\mathbf{r}). \quad (6)$$

- (c) Evaluate the force exerted by the charge q_a on another charge q_b , at position \mathbf{r}_b , using the expression for electric field you obtained using Eq. (6) in

$$\mathbf{F} = q_b \mathbf{E}(\mathbf{r}_b). \quad (7)$$

To provide a check for your calculation, the answer for the expression for the force is provided here:

$$\mathbf{F} = \frac{q_a q_b}{4\pi\epsilon_0} \frac{\mathbf{r}_b - \mathbf{r}_a}{|\mathbf{r}_b - \mathbf{r}_a|^3}. \quad (8)$$