Homework No. 08 (Spring 2015)

PHYS 520B: Electromagnetic Theory

Due date: Friday, 2015 May 8, 4.30pm

- 1. (50 points.) A particle, of charge q and mass m, always moves with speed $v \ll c$.
 - (a) Consider the case when it oscillates on the x-axis with frequency ω_0 and amplitude A given by

$$\mathbf{r}_1(t) = \hat{\mathbf{x}} A \cos \omega_0 t. \tag{1}$$

Obtain expressions for the radiated electric field $\mathbf{E}(\mathbf{r}, t)$, radiated magnetic field $\mathbf{B}(\mathbf{r}, t)$, angular distribution of the radiated power $dP/d\Omega$, and the total power radiated P.

(b) Next, consider the case when the particle moves on a circle described by

$$\mathbf{r}_2(t) = \hat{\mathbf{x}}A\cos\omega_0 t + \hat{\mathbf{y}}A\sin\omega_0 t.$$
(2)

Obtain expressions for the radiated electric field $\mathbf{E}(\mathbf{r}, t)$, radiated magnetic field $\mathbf{B}(\mathbf{r}, t)$, angular distribution of the radiated power $dP/d\Omega$, and the total power radiated P.

- (c) Show that the radiated electric and magnetic field is additive, that is, it is the sum of two oscillators.
- (d) Show that the radiated power is not additive, but exhibits interference effects. Identify the interference term for the circular motion.
- (e) Find directions $\hat{\mathbf{r}}$ for which the interference term goes to zero.
- 2. (20 points.) (Schwinger et al., problem 32.1.) A particle, of charge q and mass m, moves with speed $v \ll c$, in a uniform magnetic field **B**. Suppose the motion is confined to the plane perpendicular to **B**. Calculate the power radiated P in terms of B and v, and show that

$$P = -\frac{dE}{dt} = \gamma E,\tag{3}$$

where $E = mv^2/2$ is the energy of the particle. Find γ . Since then

$$E(t) = E(0) e^{-\gamma t},\tag{4}$$

 $1/\gamma$ is the mean lifetime of the motion. For an electron, find $1/\gamma$ in seconds for a magnetic field of $10^4\,{\rm Gauss.}$