

## Homework No. 09 (Fall 2023)

### PHYS 205A-002: UNIVERSITY PHYSICS

*School of Physics and Applied Physics, Southern Illinois University–Carbondale*

Due date: Monday, 2023 Nov 6, 2:00 PM, on D2L

### Instructions

- You are encouraged to use any of the resources to complete this homework. However, the extent to which you depend on resources while doing homework is usually a measure of how much extra work you need to put in to master the associated concepts. Solutions should be the last resource.
- Describe your thought process in detail and organize it clearly. Make sure your answer has units and the right number of significant digits.
- After completion, scan the pages as a single PDF file, and submit the file on D2L (under Assessments → Assignments).

### Problems

1. (**10 points.**) A ball having a mass of 150 g strikes a wall with a speed of 5.0 m/s and rebounds with only 50 % of its initial kinetic energy.
  - (a) What is the speed of the ball immediately after rebounding?
  - (b) If the ball was in contact with the wall for for 8.0 ms, what was the magnitude of the average force on the ball from the wall during this time interval?

[[Solution](#)]

2. (**10 points.**) A shooter of mass 90.0 kg shoots a bullet of mass 3.00 g in a direction  $60.0^\circ$  with respect to the horizontal, standing on a frictionless surface at rest. If the muzzle velocity of the bullet is 600.0 m/s, what is the recoil speed of the shooter?

[[Solution](#), see [2018S MT-03 P05](#), [2014F MT-03 P04](#)]

3. (**10 points.**) A car of mass  $m_1 = 2000.0$  kg is moving at speed  $v_{1i} = 35.0$  m/s towards East. A truck of mass  $m_2 = 5000.0$  kg is moving at speed  $v_{2i} = 25.0$  m/s towards South. They collide at an intersection and get entangled (complete inelastic collision). What is the magnitude and direction of the final velocity of the entangled automobiles?

[[Solution](#), see [2023S MT-03 P06](#), [2022F MT-03 P07](#), [2021S MT-03 P07](#), [2016F MT-03 P05](#)]




Figure 1: Problem 4.

4. (10 points.) Two masses,  $m_1 = 1.0$  kg and  $m_2 = 2.0$  kg are hanging off separate strings. Forst mass  $m_1$  is pulled to a height  $h_1 = 1.0$  m and dropped. It swings down and collides with the other hanging mass ( $m_2$  at rest) and they stick to each other (complete inelastic collision). See Figure 1. The collision happens in a plane. How high do the masses rise together after the collision.

[Solution]

5. (10 points.) What is the ratio of the final kinetic energy to initial kinetic energy in a perfectly inelastic collision involving two particles of masses  $m$  and  $M$  when the mass  $M$  is initially at rest? Express your answer in terms of  $m$  and  $M$ .

[Solution]

6. (10 points.) A mass  $m_1 = 100.$  kg moving with a speed  $v_{1i} = +10.$  m/s (elastically) collides with another mass  $m_2 = 1.0$  kg initially at rest. Determine the magnitude and direction of the final velocities of the masses after collision.

[Solution, see 2022S MT-03 P06, 2021S MT-03 P08, 2016F MT-03 P06, 2015F MT-03 P07]

7. (10 points.) Consider a thin rod of length  $L$  placed on the positive  $x$ -axis with one end at the origin. It has a mass density described by

$$\rho(x) = a + bx + cx^2, \quad a = 0 \quad b = 1.0 \frac{\text{kg}}{\text{m}^2}, \quad c = -0.8 \frac{\text{kg}}{\text{m}^3}, \quad (1)$$

where  $x$  is the distance from end placed at the origin. At what distance (in terms of  $L$ ) from the end placed at the origin is the center of of mass of the rod?

[Solution, see 2022S MT-03 P07, 2017F MT-03 P06, 2016F MT-03 P08, 2015F MT-03 P08, 2014F MT-03 P08]