

## (Preview of) Final Exam (2024 Spring)

### PHYS 510: CLASSICAL MECHANICS

*School of Physics and Applied Physics, Southern Illinois University–Carbondale*

Date: 2024 May 9

1. **(20 points.)** Not available in preview mode.
2. **(20 points.)** Not available in preview mode.
3. **(20 points.)** The path of a relativistic particle moving along a straight line with constant (proper) acceleration  $\alpha$  is described by equation of a hyperbola

$$z^2 - c^2t^2 = z_0^2, \quad z_0 = \frac{c^2}{\alpha}. \quad (1)$$




Figure 1: Problem 3

- (a) This represents the world-line of a particle thrown from  $z > z_0$  at  $t < 0$  towards  $z = z_0$  in region of constant (proper) acceleration  $\alpha$  as described by the bold (blue) curve in the space-time diagram in Figure 3. In contrast a Newtonian particle moving with constant acceleration  $\alpha$  is described by equation of a parabola

$$z - z_0 = \frac{1}{2}\alpha t^2 \quad (2)$$

as described by the dashed (red) curve in the space-time diagram in Figure 3. Show that the hyperbolic curve

$$z = z_0 \sqrt{1 + \frac{c^2 t^2}{z_0^2}} \quad (3)$$

in regions that satisfy

$$t \ll \frac{c}{\alpha} \quad (4)$$

is approximately the parabolic curve

$$z = z_0 + \frac{1}{2} \alpha t^2 + \dots \quad (5)$$

- (b) Recognize that the proper acceleration  $\alpha$  does not have an upper bound.
- (c) A large acceleration is achieved by taking an above turn while moving very fast. Thus, turning around while moving close to the speed of light  $c$  should achieve the highest acceleration. Show that  $\alpha \rightarrow \infty$  corresponding to  $z_0 \rightarrow 0$  represents this scenario. What is the equation of motion of a particle moving with infinite proper acceleration. To gain insight, plot world-lines of particles moving with  $\alpha = c^2/z_0$ ,  $\alpha = 10c^2/z_0$ , and  $\alpha = 100c^2/z_0$ .
4. (20 points.) A relativistic particle in a uniform magnetic field is described by the equations

$$\frac{dE}{dt} = \mathbf{F} \cdot \mathbf{v}, \quad (6a)$$

$$\frac{d\mathbf{p}}{dt} = \mathbf{F}, \quad (6b)$$

where

$$E = mc^2 \gamma, \quad (7a)$$

$$\mathbf{p} = m\mathbf{v}\gamma, \quad (7b)$$

and

$$\mathbf{F} = q\mathbf{v} \times \mathbf{B}. \quad (8)$$

Show that

$$\frac{d\gamma}{dt} = 0. \quad (9)$$

Then, derive

$$\frac{d\mathbf{v}}{dt} = \mathbf{v} \times \boldsymbol{\omega}_c, \quad (10)$$

where

$$\boldsymbol{\omega}_c = \frac{q\mathbf{B}}{m\gamma}. \quad (11)$$

Compare this relativistic motion to the associated non-relativistic motion.

5. (20 points.) A relativistic particle in a uniform electric field is described by the equations

$$\frac{dE}{dt} = \mathbf{F} \cdot \mathbf{v}, \quad (12a)$$

$$\frac{d\mathbf{p}}{dt} = \mathbf{F}, \quad (12b)$$

where

$$E = mc^2\gamma, \quad (13a)$$

$$\mathbf{p} = m\mathbf{v}\gamma, \quad (13b)$$

and

$$\mathbf{F} = q\mathbf{E}. \quad (14)$$

Let us consider the configuration with the electric field in the  $\hat{y}$  direction,

$$\mathbf{E} = E\hat{y}, \quad (15)$$

and initial conditions

$$\mathbf{v}(0) = 0\hat{x} + 0\hat{y} + 0\hat{z}, \quad (16a)$$

$$\mathbf{x}(0) = 0\hat{x} + y_0\hat{y} + 0\hat{z}. \quad (16b)$$

(a) In terms of the definition

$$\boldsymbol{\omega}_0 = \frac{1}{c} \frac{q\mathbf{E}}{m}, \quad (17)$$

show that the equations of motion are given by

$$\frac{d\gamma}{dt} = \boldsymbol{\omega}_0 \cdot \boldsymbol{\beta} \quad (18)$$

and

$$\frac{d}{dt}(\boldsymbol{\beta}\gamma) = \boldsymbol{\omega}_0. \quad (19)$$

(b) Since the particle starts from rest show that we have

$$\boldsymbol{\beta}\gamma = \boldsymbol{\omega}_0 t. \quad (20)$$

For our configuration this implies

$$\beta_x = 0, \quad (21a)$$

$$\beta_y\gamma = \omega_0 t, \quad (21b)$$

$$\beta_z = 0. \quad (21c)$$

Further, deduce

$$\beta_y = \frac{\omega_0 t}{\sqrt{1 + \omega_0^2 t^2}}. \quad (22)$$

Integrate again and use the initial condition to show that the motion is described by

$$y - y_0 = \frac{c}{\bar{\omega}_0} \left[ \sqrt{1 + \bar{\omega}_0^2 t^2} - 1 \right]. \quad (23)$$

Rewrite the solution in the form

$$\left( y - y_0 + \frac{c}{\omega_0} \right)^2 - c^2 t^2 = \frac{c^2}{\omega_0^2}. \quad (24)$$

This represents a hyperbola passing through  $y = y_0$  at  $t = 0$ . If we choose the initial position  $y_0 = c/\omega_0$  we have

$$y^2 - c^2 t^2 = y_0^2. \quad (25)$$

(c) The (constant) proper acceleration associated with this motion is

$$\alpha = \omega_0 c = \frac{c^2}{y_0}. \quad (26)$$

A Newtonian particle moving with constant acceleration  $\alpha$  is described by equation of a parabola

$$y - y_0 = \frac{1}{2} \alpha t^2. \quad (27)$$

Show that the hyperbolic curve

$$y = y_0 \sqrt{1 + \frac{c^2 t^2}{y_0^2}} \quad (28)$$

in regions that satisfy

$$\omega_0 t \ll 1 \quad (29)$$

is approximately the parabolic curve

$$y = y_0 + \frac{1}{2} \alpha t^2 + \dots \quad (30)$$